

Prop: $\overline{PL[0,1]} = C[0,1]$

Pf: Let $f \in C[0,1]$ and let $\epsilon > 0$.


Job: Find $g \in PL[0,1]$ s.t. $\|g - f\|_\infty < \epsilon$.

Since $[0,1]$ is compact, f is uniformly continuous so we

can find $n \in \mathbb{N}$ such that if $a, b \in [0,1]$ and $|b-a| \leq \frac{1}{n}$


then $|f(b) - f(a)| < \epsilon$.

Let $x_k = \frac{k}{n}$ $0 \leq k \leq n$.



Let g be the piecewise linear function that

equals f at each x_k and that is linear on each $[x_{k-1}, x_k]$.



Observe that on $[x_{k-1}, x_k]$

$$g(x) = f(x_{k-1}) +$$

$$\frac{x - x_{k-1}}{x_k - x_{k-1}} (f(x_k) - f(x_{k-1}))$$

Let $x \in [0, 1]$ and pick k so that $x \in [x_{k-1}, x_k]$.

Then

$$|f(x) - g(x)| = \left| f(x) - \left[f(x_{k-1}) + \frac{x - x_{k-1}}{x_k - x_{k-1}} (f(x_k) - f(x_{k-1})) \right] \right|$$

$$\leq |f(x) - f(x_{k-1})| + \left| \frac{x - x_{k-1}}{x_k - x_{k-1}} \right| |f(x_k) - f(x_{k-1})|$$

$$\leq \varepsilon + 1 \varepsilon$$

$$= 2\varepsilon.$$

Hence $\|f - g\|_0 \leq 2\varepsilon$, $\square$

We showed that PL functions with partitions at rational endpoints are dense,

Exercise: show that the PL functions as _____
with rational values at the partition points are dense.

Consequence: $C[0,1]$ is separable. $\left(c_0 \leq l_\infty \right)$

Exercise: Show that $B[0,1]$ (bounded functions on $[0,1]$)
is not separable.

Lemma: Let X be a normed vector space and W a subspace.
Then $\overline{W}$ is a subspace.

Pf: Let $a, b \in \overline{W}$. There exist sequences (a_n) and (b_n) in W converging to a, b . Then $(a_n + b_n)$ is a sequence in W and $a_n + b_n \rightarrow a + b$.

Hence $a + b \in \overline{W}$.


Scalar multiplication is handled similarly,

Hard fact:

Prop: Given $\epsilon > 0$ there exists a polynomial p such that

$$| \text{abs}(x) - p(x) | < \epsilon \text{ for all } x \in [-1, 1].$$

$$(\text{abs} \in \overline{P[-1, 1]}).$$




We'll show that

$$P[0, 1] \subseteq \overline{P[0, 1]}$$

$$\text{abs}_a(x) = \text{abs}(x-a)$$

$$[0,1]$$

$$0 \leq a \leq 1$$



$$\text{abs}_a \in \overline{P[0,1]}$$

Let $\epsilon > 0$. Find $q \in P[-1,1]$ with $\|\text{abs} - q\|_\infty < \epsilon$.


↖ on $[-1,1]$

Then if $x \in [0,1]$ then $x-a \in [-1,1]$

$$\text{So } |\text{abs}(x-a) - q(x-a)| < \epsilon$$

$$\text{i.e. } |\text{abs}_a(x) - p(x)| < \epsilon \quad \text{for all } x \in [0,1].$$

$$\| \text{abs}_a - p \|_\infty < \epsilon_0$$




$$R_a \in \overline{P[0, 1]}$$

$\uparrow$
 linear combo
 of things in
 $\overline{P[0, 1]}$

$$R_a(x) = \frac{1}{2} ((x-a) + \text{abs}_a(x))$$

$$\text{abs}_a(x) = \begin{cases} x-a & x \geq a \\ -(x-a) & x < a \end{cases}$$




$$0 \leq a < b \leq 1$$

$$V_{a,b} = \frac{R_a - R_b}{b - a}$$

$$V_{a,b} \in \overline{P[0,1]}$$

"hat"

$$a < b < c$$




$$H_{a,b,c}$$

$$H_{a,b,c} = V_{a,b} - V_{b,c}$$

$H_{a,b,c} \in \overline{P[0,1]}$ since $V_{a,b}$ and $V_{b,c}$ are

$$f \in PL[0,1], \quad f(0)=0, \quad f(1)=0$$



$$\sum_{k=1}^{n-1} y_k H_{x_{k-1}, x_k, x_{k+1}}(x) = f(x)$$

$\uparrow$ linear combo of hat functions,
 so is in $\overline{P[0,1]}$

Any $f \in PL[0,1]$ is a linear combination of
a linear function and a PL function that vanishes
at 0 and 1.

Prop: Given $\varepsilon > 0$ there exists a polynomial $p(x)$ with
 $| \text{abs}(x) - p(x) | < \varepsilon$ for all $x \in [-1, 1]$.

Pf: Let $P_0(x) = 0$ and if P_n has been defined,
let
$$P_{n+1}(x) = P_n(x) + \frac{1}{2} (x - P_n(x))^2$$

Claim $P_n \rightarrow \text{sgn}$ uniformly on $[0,1]$.

We first show that for all n ,

$$P_{n+1} \geq P_n \geq 0$$

and that $\sqrt{x} \geq P_n(x) \geq 0$ for all $x \in [0,1]$.

These facts are clear if $n=0$.

Suppose these facts hold for some n .

Observe that

$$P_{n+1}(x) = P_n(x) + \frac{1}{2} (\sqrt{x} + P_n(x)) \cdot (\sqrt{x} - P_n(x)).$$

Moreover $\frac{1}{2}(\sqrt{x} + P_n(x)) \leq \frac{1}{2}(\sqrt{x} + \sqrt{x}) \leq 1$

and $\sqrt{x} - P_n(x) \geq 0$.

$$\text{Hence } P_{n+1}(x) \leq P_n(x) + 1 \cdot (\sqrt{x} - P_n(x)) \\ = \sqrt{x}.$$

$$\text{Moreover, } P_{n+1}(x) = P_n(x) + \frac{1}{2} (x - P_n(x)^2) \\ \geq P_n(x) \geq 0.$$

Finally, $P_{n+2}(x) \geq P_{n+1}(x)$ for the same reasons.

Now for each $x \in [0, 1]$, $P_n(x)$ is a monotone increasing sequence in $\mathbb{R}$ that is bounded above by $\sqrt{x}$. So, $\{P_n\}$ converges to some $f(x)$.

That is $P_n \rightarrow f$ pointwise on $[0, 1]$.

Using

$$P_{n+1}(x) = P_n(x) + \frac{1}{2}(x - P_n(x))^2$$

we take a limit and conclude

$$f(x) \equiv f(x) + \frac{1}{2}(x - f(x))^2$$

so $f(x)^2 = x$. But $f(x) \geq 0$, so $f(x) = \sqrt{x}$.

Since $\sqrt{x}$ is continuous and since P_n is monotone increasing in n , $P_n \rightarrow f$ uniformly.