

Last class: compactness.

Def: Let X be a metric space. A set $A \subseteq X$ is bounded if there exists $x \in X$ and $r > 0$ such that $A \subseteq B_r(x)$.

Note: there is complete flexibility in x . (Δ inequality)

Lemma: Compact subsets of a metric space are bounded.

Pf: Suppose $A \subseteq X$ is not bounded.

Pick $x \in X$ and form the cover $\{B_r(x)\}_{r>0}$

These sets cover A but because A is not bounded

there is no finite subcover, so A is not compact.

(Cor: In metric spaces compact sets are closed and bounded.

The converse (even for metric spaces) is false.

e.g. $\mathbb{Z}$ $d(x, y) = 1 \quad x \neq y,$

Is $\mathbb{Z}$ bounded? Yep!

Is $\mathbb{Z}$ closed in $\mathbb{Z}$? Yep!

Is $\mathbb{Z}$ compact? No: the open cover by singletons has no finite subcover.

Next semester: compact $\Leftrightarrow$ complete + totally bounded in metric spaces

But! In $\mathbb{R}^n$, compact $\Leftrightarrow$ closed + bounded holds.

(Extreme Value Theorem) Using: subsets of $\mathbb{R}$ are compact $\Leftrightarrow$ closed + bounded - and f is
Prop: Suppose $f: K \rightarrow \mathbb{R}$ where K is compact and nonempty, cto.

Then there exist $k_-, k_+ \in K$ such that

$$f(k_-) \leq f(k) \leq f(k_+) \quad \text{for all } k \in K.$$

Pf: Observe $f(K) \subseteq \mathbb{R}$ is compact. Then $f(K)$ is closed and bounded. Since it is nonempty (as K is nonempty) and bounded above, it admits a supremum s . There is a sequence in $f(K)$ converging to s and hence

$s \in \overline{f(K)} = f(K)$. Hence there is $k_+ \in K$ with $f(k_+) = s$.

So $f(k_+) \geq f(k)$ for all $k \in K$ as s is an upper bound for $f(K)$. The construction of k_- is similar.

$A \subseteq \mathbb{R}$, closed, bounded

bounded $\Rightarrow A \subseteq [-R, R]$ for some R .

$[-R, R] \sim [0, 1]$ which is compact.

A is closed in $\mathbb{R}$ and contained in $[-R, R]$
and is hence closed in $[-R, R]$

Hence A is compact as a subspace of $[-R, R]$
but then also as a subspace of $\mathbb{R}$.

How to generalize this reasoning to $\mathbb{R}^n$?

Steps: 1) $[-R, R]$ is compact for all $R > 0$.

2) If $X_1, \dots, X_n$ are compact then

$X_1 \times \dots \times X_n$ is compact.

3) $[-R, R]^n$ is compact w/ product top
but then also as a subspace of $\mathbb{R}^n$.

4) If $A \subseteq \mathbb{R}^n$ is bounded then
 $A \subseteq [-R, R]^n$ for some large R .

$$B_r(0) \quad x = (x_1, \dots, x_n) \\ \hookrightarrow \subseteq [-r, r]^n.$$

5) If in addition A is closed then
the above argument shows A is compact.



Lemma: (Tube Lemma)

Suppose X, Y are spaces and Y is compact.
 Let $x_0 \in X$ and let W be an open set
 containing $\{x_0\} \times Y$. Then there is an open
 set $U \subseteq X$ with $x_0 \in U$ and

$$\underbrace{U \times Y}_{\hookrightarrow \pi_X^{-1}(U)} \subseteq W.$$

Pf: For each (x_0, y) we can find a basic
 open set $U_y \times V_y \subseteq W$ with

$x_0 \in U_y$ and $y \in V_y$. The fiber $\{x_0\} \times Y$
 is homeomorphic to Y and hence compact.



Then sets $U_\gamma \times V_\gamma$ cover the fiber,


so we can find a finite subcover $U_{\gamma_1} \times V_{\gamma_1}, \dots, U_{\gamma_n} \times V_{\gamma_n}$.


Let $U = \bigcap_{j=1}^n U_{\gamma_j}$. Then U is open in X and contains x_0 .

Suppose $(x, y) \in U \times Y$. Pick j with $(x_0, y) \in U_{\gamma_j} \times V_{\gamma_j}$.

Then $(x, y) \in U \times V_{\gamma_j} \subseteq U_{\gamma_j} \times V_{\gamma_j} \subseteq W$.

So $U \times Y \subseteq W$.





$$\{U_\alpha\}$$

$\{x\} \times Y$ is covered by finitely many U_α 's.

$$\pi_x^{-1}(\{x\}) \subseteq \bigcup_{\alpha \in A_x} U_\alpha \text{ where } A_x \text{ is finite.}$$

$$W_x = \bigcup_{\alpha \in A_x} U_\alpha \text{ is open and contains } \{x\} \times Y.$$

$$\text{Find } U_x \text{ with } U_x \times Y \subseteq W_x$$

The open sets U_x cover X , so we can extract a finite subcover, $U_{x_1}, \dots, U_{x_n}$.

$$X \subseteq \bigcup U_{x_j} \quad \text{and} \quad X_{x_j} \subseteq \bigcup \pi_{x_j}^{-1}(U_{x_j})$$

Each $\pi_{x_j}^{-1}(U_{x_j})$ is contained in W_{x_j} and hence covered by finitely many sets U_α . But there are finitely many tubes $\pi_{x_j}^{-1}(U_{x_j})$.

How about arbitrary products of compact sets?

This is false in the box topology.

$\underbrace{\{0, 1\}^\omega}_{\text{discrete!}}$ is discrete with the box topology.

discrete!

The singletons are an open cover w/ no finite subcover.

For the product topology this is true! (Tychonoff's Thm).