

Prop: If X is limit point compact and 1^{st} countable and Hausdorff
then it is sequentially compact.

Pf: Let $\{x_n\}$ be a sequence in X . If the sequence takes on only finitely many values then we can extract a constant subsequence. Hence we assume the sequence has infinitely many values and these have a limit point x_0 .

n_k
↑

If x is an element of the sequence we reduce to a subsequence where none of the terms are x but still we take on infinitely many values.



Let $\{B_j\}$ be a countable nested neighborhood base at x .

Pick n_1 such that $x_{n_1} \in B_1$. This is possible since x is a limit point of the sequence. Now suppose we have selected $n_1 < n_2 < \dots < n_k$ such that $x_{n_j} \in B_j$.

Notice that $B_{k+1} \setminus \{x_1, x_2, \dots, x_{n_k}\}$ is open (Hausdorff!)

and contains x . Hence it contains a term of the sequence

$x_{n_{k+1}}$ with $n_{k+1} > n_k$. Inductively we have obtained a

subsequence with $x_{n_j} \in B_j$ for all j .

Now let U be an open set containing x .

There exists J with $x \in B_J \subseteq U$.

If $j \geq J$ then $x_{n_j} \in B_j \subseteq B_J \subseteq U$.

Hence $x_{n_j} \rightarrow x$.

$\mathcal{U}(x)$
 $\uparrow$
all open sets
containing x .



Prop: A sequentially compact 2^{nd} countable space X is compact.

Pf: Let $\{U_\alpha\}_{\alpha \in I}$ be an open cover of X .

Since X is 2^{nd} countable it is Lindelöf and we can extract a countable subcover, call it $\{U_j\}$.

Suppose to the contrary that there is no finite subcover.

For each j we can then pick $x_j \in X$, $x_j \notin U_1 \cup \dots \cup U_j$.

The space is sequentially compact, so we can find a convergent subsequence x_{j_k} converging to some x .

Pick J such that $x \in U_J$. Then there exists K

so that if $k \geq K$, $x_{j_k} \in U_J$. But $x_{j_k} \notin U_J$

if $j_k \geq J$. So if $k \geq K$ and $j_k \geq J$ (which

we can assure by taking k large enough) $x_{j_k} \in U_J$ and $x_{j_k} \notin U_J$, a contradiction.

$\mathbb{R}P^2$ is metrizable but no particular metric seems important.

For metric spaces The seq compact $\Rightarrow$ compact result boils down to a question of separability.

(i.e. whether the space has a countable dense subset.)

Suppose X is a separable metric space. Then it is 2^{nd} countable. So if it is sequentially compact then it is compact.

We'll show shortly that a sequentially compact metric space is in fact separable!

Here, by the previous argument, a sequentially compact metric space is compact. Moreover, if a metric space is not separable then it is not sequentially compact. so it is not compact.

$$\text{compact} \Rightarrow \text{l.p.c} \Rightarrow \text{seq. compact}$$

Key prop: in a sequentially compact metric space,

for all $\varepsilon > 0$ there is a covering by finitely many ε -balls

↳ ball of radius ε .

To get separability: For each $\varepsilon = \frac{1}{n}$ find finitely many x_n 's such that

$$\bigcup_j B_{\frac{1}{n}}(x_{n,j}) = X.$$

Countably many $x_{n,j}$'s and they are dense.

Lemma: Let X be a sequentially compact metric space.

For all $\varepsilon > 0$ there exists a cover of X by finitely many ε -balls.

Pf: Let $\varepsilon > 0$ and suppose to the contrary that no

such cover exists. Pick $x_1 \in X$. Since $B_\varepsilon(x_1)$ is not a cover we can find $x_2 \notin B_\varepsilon(x_1)$. Since $\{B_\varepsilon(x_j)\}_{j=1}^2$

is not a cover we can find x_3 such that

$d(x_3, x_1) \geq \varepsilon$ and $d(x_3, x_2) \geq \varepsilon$. Continuing inductively

we can find a sequence $\{x_j\}$ with $d(x_j, x_l) \geq \varepsilon$ if $j \neq l$. The space is sequentially compact and

we have a convergent subsequence $x_{j_k} \rightarrow x$.
Hence we can find K so that if $k \geq K$,

$d(x_{j_k}, x) < \frac{\varepsilon}{10}$. But then if $k_1, k_2 \geq K$,

$$d(x_{j_{k_1}}, x_{j_{k_2}}) \leq \frac{2\varepsilon}{10} = \frac{\varepsilon}{5} \text{ by the } \bigcirc$$

Δ inequality. But $d(x_{j_{k_1}}, x_{j_{k_2}}) \geq \varepsilon$ if $k_1 \neq k_2$, a contradiction.